

## EFFECT OF BIOCHEMICAL PARAMETERS ON MALARIA INFECTED PATIENTS IN GENERAL HOSPITAL BILLIRI, GOMBE STATE NIGERIA

<sup>1</sup>PHILIMON J, <sup>2</sup>CHESSSED G, <sup>2</sup>PUKUMA, <sup>1</sup>ABBA E., <sup>3</sup>AMINA Y.U., <sup>1</sup>ISHAKU M., <sup>1</sup>LAMOGO Y.,  
<sup>1</sup>JEMIMAH <sup>4</sup>A., SARKI <sup>5</sup>A. MUHAMMAD A.A., <sup>5</sup>SHUAIBU S. and <sup>1</sup>FATAI, A.

<sup>1</sup>Department of Zoology, Faculty of Science, Gombe state University, Gombe state Nigeria

<sup>2</sup>Department of Zoology, School of Life Sciences, Modibbo Adama University of Technology Yola,  
Adamawa State Nigeria

<sup>3</sup>Department of Biological Sciences, Faculty of Science, Abubakar Tafawa Balawa University, Bauchi  
State

<sup>4</sup>Department of Biological Sciences, Faculty of Science, Federal University Kashere, Gombe State

<sup>5</sup>Department of Biology, School of Secondary School Education (Science), Federal College of Education  
Technical Gombe, Gombe State Nigeria

**Corresponding email:** philimonjay@gsu.edu.ng

### ABSTRACT

This study was carried out to determine the impact of malaria parasite infection caused by *Plasmodium falciparum* on some biochemical parameters. One hundred and five (105) malaria parasite positive male and female subjects and fifty (50) apparently healthy subjects were recruited for this study. Malaria Parasite diagnosis was done by Microscopic Examination of Thick and Thin Stained Blood Films. Age range 15-25 had the highest prevalent rate of 17.14% for males and 14.29% for females while the age range 55-60 had the least prevalent rate of 0.95% for both male and female. when control values were compared with the mean values obtained from the biochemical analysis of *Plasmodium falciparum* parasitaemia subject samples, it was significantly lower for glucose ( $p=0.04$ ), ALT ( $p=0.00$ ), AST ( $p=0.00$ ), ALP ( $p=0.00$ ), total protein ( $p=0.01$ ), Globulin ( $p=0.02$ ) and Creatinine ( $p=0.04$ ). Total cholesterol ( $p=0.12$ ), Triglyceride ( $p=0.06$ ), Albumin ( $p=0.49$ ) and Urea ( $p=0.15$ ) were statistically not significant for malaria parasitaemia subject samples. Glucose, Total cholesterol, Liver enzymes, creatinine changes were seen involved in malaria infection caused by *Plasmodium falciparum*. The prevention of malarial infection is still the most valid method of preventing these conditions and early diagnosis and treatment are the measures likely to decrease malarial complications commonly determined by changes in these biochemical parameters.

**Keywords:** Malaria, Parasitaemia, Biochemical Profile, Infected, *P. falciparum*

### INTRODUCTION

Malaria parasites are members of the genus plasmodium (Phylum Apicomplex). In humans malaria is caused by *P. falciparum*, *P. malarie*, *P. ovale*, *P. vivax* and *P. knowlesi* (Singh 2015; Mueller *et al.*, 2019). Malaria is a mosquito-borne infectious disease of human caused by eukaryotic protists of the genus plasmodium. It is widespread in tropical and subtropical regions, including much of sub-saharan Africa, Asia and the America. Malaria is prevalent in these regions because of the significant amounts of rainfall and consistent high humidity along with stagnant water in which their larvae mature and provide mosquitoes in the environment (Prothero and Mansell, 2020). Malaria was once common in most of Europe and North America (Lindemann, 2018) where it is no longer endemic (Gratz, 2017) though imported cases do occur. A total number of 216 million estimated malaria cases occurred in 2008 as reported by World Health Organization. A total of

106 countries of the world were highly affected of which Nigeria was included (World Health Organization, 2021) 81% of malaria cases occurred in the Africa region and 43% of the reported cases occurred in Nigeria out of which 655,000 deaths was reported worldwide. Malaria develops via two phases: an exoerythrocytic and an erythrocytic phase. The exoerythrocytic phase involves infection of the hepatic system or liver, whereas the erythrocytic phase involves infection of the erythrocyte or red blood cell when an infected female anopheles mosquito pierces a person's skin to take a blood meal, sporozoites in the mosquito saliva enters the blood stream and migrate to the liver. Within minutes of being introduced into the human host, the sporozoites infect hepatocytes, multiplying asexually and asymptotically for a period of 8-30 days (Bledsoe, 2017). Once in the liver, these organisms differentiate to yield thousands of merozoites, which following rupture of their host cells, escape into the blood and infect red blood

cells, thus beginning the erythrocytic stage of the life cycle (Bledsoe, 2017). The parasite escapes from the liver undetected by wrapping itself in the cell membrane of the infected host liver cell (Sturn, 2019). Within the red blood cells, the parasite multiply further, again asexually, periodically breaking out of their hosts to invade fresh blood cells. Several such amplification cycles occur (Sturn, 2019). Thus classical description of waves of fever arises from simultaneous waves of merozoites escaping and infecting red blood cells. Some *P. vivax* and *P. ovale* sporozoites do not immediately develop into exoerythrocytic phase merozoites but instead produce hypozoites that remain dormant for period ranging from several months (6-12 months is typical) to as long as three

years. After a period of dormancy, they reactivate and produce merozoites. Hypozoites are responsible for long incubation and late relapses in these two species of malaria (Cogswell, 2019). It was hypothesized that the malaria parasite uses cholesterol and phospholipids from its host, resulting in a decrease of serum HDL. Prior to this report, Angus *et al.* (2015). utilized lipoprotein electrophoresis in rhesus monkeys infected with *Plasmodium knowlesi* to study serum lipids in malaria. Their results were not conclusive because lipoprotein bands could barely be detected in the serum of controls. Subsequently, several clinical studies showed lipid profile changes in the setting of both uncomplicated and complicated malaria (Angus *et al.*, 2015 and Chagnon *et al.*, 2017).

## Materials and Methods

### Research Design

A total of one hundred and five (105) malaria parasite positive male and female subjects between the ages of 15-60 years visiting general hospital laboratories in Billiri town and adjoining rural settlements were recruited for this study. A total of fifty (50) (25 males and 25 females) apparently healthy male and female subjects between the ages of 15-60 years who tested negative for malaria parasite were recruited as control subjects for this study. Ethical approval was obtained and the study was conducted with informed consent of the patients.

### Geographical Description of the Study Area

This study was carried out in Billiri town in Gombe state, Nigeria. It is a semi urban area with an estimated population of 125,842 people at the 2006 census. (National Population Commission 2006).

### Sample Collection

Ten mililitre (10ml) of venous blood samples were collected by vene-puncture from the subjects into accurately labeled plain containers for both subjects and control. For this study, the test subjects were those that presented at the various health centres and General hospitals with symptoms associated with malaria infection. The signs and symptoms of malaria infection in humans are caused by the asexual blood stage of

the parasite which includes: headache, joint pains, nausea, vomiting, abdominal upset, fever, and digestive disorders (Dayaci, 2018). Thick and thin stained blood films were made, and the remaining blood samples were centrifuged at 3500 rpm for ten minutes at room temperature within one hour of sample collection and the serum separated into clean plain containers which were kept frozen until required for analysis.

### Malaria Parasite Diagnosis by Microscopic Examination of Thick and Thin Stained Blood Films

The Giemsa's staining technique was used for the staining of the thick blood films for malaria parasite detection and malaria parasite count, while the thin blood films were stained with Leishman staining technique for plasmodium species identification as described by Monica Cheesbrough (2005). The changes in parasitized red cells helped to identify plasmodium species and to detect mixed infection of malaria parasite. The number of asexual *Plasmodium falciparum* and other species per 200 leukocytes were counted and when ten or more parasites were identified, the number was recorded, a blood sample was regarded as negative when the examination of thick films failed to show the presence of asexual parasites. The parasite count in relation to the leukocyte count was converted to parasite per micro litre of blood using this mathematical formula;

$$\frac{\text{Number of parasites}}{\text{Number of leukocytes}} \times \text{Total WBC count} = \text{Parasite per micro litre of blood}$$

### Procedure for Staining Thick Blood Film

This involved making thick blood films on clean

grease free glass slides, allowed to air dry and stained with prepared Giemsa stain for 30 minutes.

The Giemsa stain was prepared by diluting stock Giemsa stain (commercially obtained) in buffered water immediately before use. Stained slides were rinsed in clean water and allowed to air dry before examination under a microscope using X100 objective lens. Chromatin of malaria parasite was stained dark red and cytoplasm stained blue with Giemsa's stain. The presence of malaria parasite, identification of the species of human parasites and relative malaria parasite count in each blood sample was determined from the Giemsa stained thick films and Leishman stained thin blood films. Malaria Parasitaemia was confirmed by microscopic examination using X100 objective lens (oil immersion lens). A slide was scored as negative when 100 high power fields had been examined for about 30 minutes without seeing any parasites. The amount of relative parasite count (Occurrence) in positive smears was done using a simple code from one to four crosses (+ - ++++), although none of the subjects had ++++. Malaria Parasitaemia Occurrence was graded as; + = 1 – 10 parasites per 100 thick film field; ++ = 11 – 100 parasites per 100 thick film field; +++ = 1 – 10 parasites per single thick film field; ++++ = more than 10 parasites per single thick film field after staining for 30 minutes as described by Monica Cheesbrough (2005).

### Procedure for Staining Thin Blood Film

Thin blood films were made on clean grease free glass slide and stained using Leishman staining technique as described by Monica Cheesbrough (2005). The films were allowed to air dry and covered with Leishman stain for four minutes. The stain on the slides were diluted with buffered distilled water and allowed to stain for ten minutes. Slides were rinsed with water, allowed to air dry and examined under microscope using X100 objective lens.

### Malaria Parasite Count

Quantitative parasitaemia count (Parasite density) was determined by counting the number of asexual parasites (trophozoites, schizonts) present in as many microscopic fields (100x) necessary to count 200 leukocytes in each thick blood film and multiplies by the total white blood cells count of each blood sample. Parasitaemia was graded as low (parasite 1000-9,999  $\mu\text{L}^{-1}$ ) and high ( $>10,000 \mu\text{L}^{-1}$ ).

### Biochemical Analysis

Alanine Amino Transferase (ALT) and Aspartate Amino Transferase (AST) activity was determined

using the method described by Rietman and Frankel (1957), while the activities of Alkaline Phosphatase (ALP) determined using the method described by Rec. Gscc (DGKC) (1972). The Bromo-cresol Green (BCG) binding method was used for the determination of albumin (Slot, 1977). 3% Sulphosalicylic acid (SSA) colorimetric method was used for the determination of Total protein. The Jaffe-Slot modified alkaline picrate colorimetric method was used for Creatinine level estimation (Slot, 1977). Urea levels were determined using the diactetylmonozine methods (Rey and Weiling, 1970). Glucose oxidase method was used for the estimation of plasma glucose. Plasma total Cholesterol was determined colorimetrically using cholesterol oxidase method as described by Alkin *et al.* (1974).

### Statistical Analysis

The data generated from the study was subjected to basic statistical measurement for mean and standard deviation, and comparisons were carried out to test for significant differences, using parametric analysis of variance (ANOVA) with the aid of the computer SPSS 20.0 windows application. Statistical significance was accepted at  $p < 0.05$ .

### Results

The statistical data obtained from the analyzed subjects and control samples are shown in Tables 1-5 below. Values from Control samples were compared with values from the test subjects, and multiple comparisons were carried out with the one-way ANOVA statistics between groups and within groups and the control and test subject values are presented as mean  $\pm$  Standard Deviation (SD).

### The Demographic Characteristic of the Study Population

showed that a total of 105 test subjects was used for this study, 59 (56.18%) were males of which 18 (17.14%) of them were within the ages of 15-25, those within the ages of; 26-35 were 16 (15.23%), 36-45 were 14 (13.33%), 46-55 were 11 (10.48%). 28 (26.60%) males had '+' occurrence of malaria parasitaemia, 24 (22.85%) had '++' occurrence of malaria parasitaemia and 7

(06.60%) males had ‘+++’ occurrence of malaria parasitaemia. A total of 46 (43.81%) female test subjects were used for this study, within the ages of 15-55. Those within the ages of 15-25 were 14 (13.33%), within the ages of 26-35 were 15 (14.29%), within the ages of 36-45 were 08 (07.62%), and those within the ages of 46-55 were 09 (08.57%). 30 (28.51%) females had ‘+’ occurrence of malaria parasitaemia, 14 (13.33%) had ‘++’ occurrence of malaria parasitaemia and 03 (02.81%) females had ‘+++’ occurrence of malaria parasitaemia.

As shown in table 2; when control values were compared with the mean values obtained from the biochemical analysis of *Plasmodium falciparum* parasitaemia subject samples, it was significantly lower for glucose (p=0.04), ALT (p=0.00), AST (p=0.00), ALP (p=0.00), total protein (p=0.01), Globulin (p=0.02) and Creatinine (p=0.04). Total cholesterol (p=0.12), Triglyceride (p=0.06), Albumin (p=0.49) and Urea (p=0.15) were statistically not significant for malaria parasitaemia subject samples.

**Table 1:** Mean  $\pm$  SD of some Biochemical parameters of *plasmodium falciparum* malaria infected inhabitants of Billiri metropolis compared with healthy controls

Parameter	Control	Malaria Infected Subjects	f-ratio	p-values
Glucose (mg/dl)	79.21 $\pm$ 11.24	64.34 $\pm$ 8.47	4.34	0.04*
Total Cholesterol (mg/dl)	154.65 $\pm$ 27.98	149.34 $\pm$ 20.77	6.28	0.12 0.06
Triglyceride (mg/dl) ALT (IU/L)	85.04 $\pm$ 25.39	81.48 $\pm$ 24.77	3.19	0.00*
AST (IU/L)	10.66 $\pm$ 5.34	24.34 $\pm$ 3.15	4.34	0.00*
ALP (mg/dl)	13.48 $\pm$ 3.43	28.50 $\pm$ 2.38	2.75	0.00*
Albumin (g/dl)	44.13 $\pm$ 13.65	64.23 $\pm$ 5.24	5.25	0.49
Total protein (g/dl)	3.76 $\pm$ 0.73	3.85 $\pm$ 0.32	3.02	0.01*
Globulin (g/dl)	7.85 $\pm$ 0.31	6.11 $\pm$ 0.35	4.50	0.02*
Creatinine (mg/dl)	4.09 $\pm$ 0.12	2.26 $\pm$ 0.23	2.34	0.04*
Urea (mg/dl)	0.68 $\pm$ 0.16	0.89 $\pm$ 0.44	3.45	0.15
	15.44 $\pm$ 1.12	17.68 $\pm$ 1.23	4.21	

KEY: \*Statistically significant (p<0.05), ALT; Alanine Amino Transferase, AST; Aspartate Amino transferase, ALP= Alkaline phosphatase, g/dl; gram per decilitre, IU/L; international unit per litre, mg/dl; milligram per decilitre, SD; standard deviation.

As shown in table 2 below; when male control subject values were compared with mean values obtained from the biochemical analysis of male *Plasmodium falciparum* parasitaemia subject samples, it was significantly lower for glucose (p=0.01), ALT (p=0.02), AST (p=0.01), ALP

(p=0.00) and Creatinine (p=0.04). Total cholesterol (p=0.06), Triglyceride (p=0.08), total protein (p=0.21), Globulin (p=0.78), Albumin (p=0.13) and Urea (p=0.22) were statistically not significant for malaria parasitaemia subject samples.

**Table 2:** Mean $\pm$ SD of some biochemical parameters of *Plasmodium falciparum* Malaria infected male inhabitants of Ekpoma metropolis compared with healthy male controls

Parameter	Male Control	Male Infected Subjects	f-ratio	p-values
Glucose (mg/dl)	80.12 $\pm$ 9.20	69.14 $\pm$ 8.47	3.36	0.01*
Total Cholesterol (mg/dl)	148.47 $\pm$ 20.28	145.34 $\pm$ 30.00	7.11	0.06 0.08
Triglyceride (mg/dl) ALT (IU/L)	89.04 $\pm$ 15.21	83.18 $\pm$ 11.01 25.74 $\pm$ 2.18	4.11	0.02*
AST (IU/L)	11.54 $\pm$ 4.30	27.43 $\pm$ 3.12	6.14	0.01*
ALP (mg/dl)	12.23 $\pm$ 3.13	69.13 $\pm$ 8.09	4.15	0.00*
Albumin (g/dl)	45.13 $\pm$ 8.55	4.89 $\pm$ 0.42	4.15	0.13 0.21
Total protein (g/dl)	4.16 $\pm$ 1.00	8.16 $\pm$ 1.05	2.09	0.78
Globulin (g/dl)	7.98 $\pm$ 1.01	3.22 $\pm$ 0.32	3.71	0.03*
Creatinine (mg/dl)	3.82 $\pm$ 0.76	0.95 $\pm$ 0.22	4.32	0.22
Urea (mg/dl)	0.79 $\pm$ 0.11	18.08 $\pm$ 0.91	5.33	
	16.17 $\pm$ 1.00		7.20	

KEY: \* Statistically significant (p<0.05), ALT; Alanine Amino Transferase, AST; Aspartate Amino transferase, ALP= Alkaline phosphatase, g/dl; gram per decilitre, IU/L; international unit per litre, mg/dl; milligram per decilitre, SD; standard deviation.

As shown in table 3 below; when female control subject values compared with mean values obtained from the biochemical analysis of female *Plasmodium falciparum* malaria parasitaemia subject samples, it was significantly lower for glucose (p=0.01), ALT (p=0.00), AST (p=0.01),

ALP (p=0.04) and Creatinine (p=0.00). Total cholesterol (p=0.09), Triglyceride (p=0.11), total protein (p=0.08), Globulin (p=0.13), Albumin (p=0.06) and Urea (p=0.27) were statistically not significant for malaria parasitaemia subject samples.



**Table 3:** Mean $\pm$ SD of some biochemical parameters of Plasmodium falciparum Malaria infected female inhabitants of Billiri metropolis compared with healthy female controls

Parameter	Female Control	Female Infected Subjects	f-ratio	p-values
Glucose (mg/dl)	72.22 $\pm$ 5.30	62.23 $\pm$ 5.14	4.12	0.01*
Total Cholesterol (mg/dl)	154.11 $\pm$ 23.00	150.34 $\pm$ 30.00	5.01	0.09 0.11
Triglyceride (mg/dl) ALT (IU/L)	86.01 $\pm$ 8.20	85.48 $\pm$ 14.00	2.05	0.00*
AST (IU/L)	10.13 $\pm$ 3.21	23.63 $\pm$ 1.29	7.06	0.01*
ALP (mg/dl)	11.13 $\pm$ 2.22	25.21 $\pm$ 4.00	9.04	0.04*
Albumin (g/dl)	36.10 $\pm$ 3.15	65.25 $\pm$ 6.11	5.26	0.06 0.08
Total protein (g/dl)	3.70 $\pm$ 0.90	3.51 $\pm$ 0.72	4.43	0.13
Globulin (g/dl)	7.67 $\pm$ 2.13	7.91 $\pm$ 2.00	6.22	0.00*
Creatinine (mg/dl)	3.97 $\pm$ 0.18	3.68 $\pm$ 0.67	3.15	0.27
Urea (mg/dl)	0.59 $\pm$ 0.06	0.80 $\pm$ 0.22	9.08	
	12.38 $\pm$ 1.08	14.09 $\pm$ 1.01	3.99	

As shown in table 4 below; When the various levels of severity of *Plasmodium falciparum* malaria parasite in the test subjects were compared and multiple comparisons carried out with the one-way ANOVA statistics between groups and within groups, Mean Glucose value for + (1 plus) malaria parasitaemia (67.25 $\pm$ 6.34) was significantly higher when compared with ++ (2 plus) Malaria parasitaemia (62.65 $\pm$ 4.33) and +++ (3 plus) Malaria parasitaemia (59.17 $\pm$ 5.21),  $p=0.04$ . Mean Total cholesterol value for + (1 plus) malaria parasitaemia (151.14 $\pm$ 1.77) was not significant in comparison with ++ (2 plus) Malaria parasitaemia, but was significantly higher in comparison with +++ (3 plus) Malaria parasitaemia (144.10 $\pm$ 24.23),  $p=0.03$ . Mean ALT levels for + (1 plus) malaria parasitaemia (18.34 $\pm$ 3.15) was not significant in comparison with

++ (2 plus) Malaria

parasitaemia, but was significantly lower in comparison with +++ (3 plus) Malaria parasitaemia (26.01 $\pm$ 1.19),  $p=0.01$ . Mean AST levels for + (1 plus) malaria parasitaemia (16.43 $\pm$ 2.08) was significantly lower in comparison with ++ (2 plus) (23.20 $\pm$ 1.12) and +++ (3 plus) (29.15 $\pm$ 2.02) malaria parasitaemia, ++ (2 plus) was significantly lower in comparison with +++ (3 plus) Malaria parasitaemia ( $p=0.02$ ). Mean Creatinine levels for + (1 plus) malaria parasitaemia (0.68 $\pm$ 0.32) was significantly lower in comparison with ++ (2 plus) (0.85 $\pm$ 0.15) and +++ (3 plus) (1.3 $\pm$ 0.62) malaria parasitaemia, ++ (2 plus) was significantly lower in comparison with creatinine levels for +++ (3 plus) Malaria parasitaemia ( $p=0.00$ ). Triglyceride, ALP, Albumin, Total protein, globulin and Urea were not statistically significant.

**Table 4:** Effect of different levels severity of Plasmodium falciparum parasitaemia on Some Biochemical Parameters.

Parameter	+ (1 plus) Malaria parasitaemia	++ (2 plus) Malaria parasitaemia	+++ (3 plus) Malaria parasitaemia	f-ratio	p-values
Glucose (mg/dl)	67.25 $\pm$ 6.34 <sup>a</sup>	62.65 $\pm$ 4.33 <sup>b</sup>	59.17 $\pm$ 5.21 <sup>b</sup>	2.69	0.04*
Total Cholesterol (mg/dl)	151.14 $\pm$ 1.77 <sup>a</sup>	148.21 $\pm$ 31.41 <sup>ab</sup>	144.10 $\pm$ 24.23 <sup>b</sup>	8.08	0.03*
Triglyceride (mg/dl) ALT (IU/L)	87.68 $\pm$ 24.77 <sup>a</sup>	87.08 $\pm$ 15.70 <sup>a</sup>	80.19 $\pm$ 30.11 <sup>a</sup>	1.94	0.07
AST (IU/L)	18.34 $\pm$ 3.15 <sup>a</sup>	21.21 $\pm$ 1.05 <sup>ab</sup>	26.01 $\pm$ 1.19 <sup>b</sup>	5.00	0.01*
ALP (mg/dl)	16.43 $\pm$ 2.08 <sup>a</sup>	23.20 $\pm$ 1.12 <sup>b</sup>	29.15 $\pm$ 2.02 <sup>c</sup>	6.25	0.02*
Albumin (g/dl)	59.21 $\pm$ 1.14 <sup>a</sup> 3.75 $\pm$ 0.02 <sup>a</sup>	64.00 $\pm$ 6.03 <sup>a</sup> 3.55 $\pm$ 0.32 <sup>a</sup>	69.41 $\pm$ 3.44 <sup>a</sup> 3.80 $\pm$ 0.14 <sup>a</sup>	3.12	0.10
Total protein (g/dl)	7.10 $\pm$ 1.05 <sup>a</sup>	7.41 $\pm$ 1.12 <sup>a</sup>	7.81 $\pm$ 1.63 <sup>a</sup>	3.11	0.21
Globulin (g/dl)	3.35 $\pm$ 0.34 <sup>a</sup>	3.86 $\pm$ 0.56 <sup>ab</sup>	4.01 $\pm$ 1.00 <sup>b</sup>	4.67	0.32
Creatinine (mg/dl)	0.68 $\pm$ 0.32 <sup>a</sup>	0.85 $\pm$ 0.15 <sup>b</sup>	1.3 $\pm$ 0.62 <sup>c</sup>	4.04	0.05
Urea (mg/dl)	14.23 $\pm$ 4.03 <sup>a</sup>	13.44 $\pm$ 2.05 <sup>a</sup>	16.08 $\pm$ 3.03 <sup>a</sup>	8.03	0.00*
				1.21	0.18

KEY: \* Statistically significant ( $p<0.05$ ), ALT; Alanine Amino Transferase, AST; Aspartate Amino transferase, ALP= Alkaline phosphatase, g/dl; gram per decilitre, IU/L; international unit per litre, mg/dl; milligram per decilitre, SD; standard deviation.

As shown in table 5 below, when the various age groups of malaria parasitaemia test subjects were compared and multiple comparisons carried out with the one-way ANOVA statistics between groups and within groups it was not significant for any of the biochemical parameters tested ( $p>0.05$ ).

## Discussion

The result of demographic characteristic of the study population showed that males had higher prevalence of Malaria parasitaemia than females, which agrees with studies done by Mbanugo and

Ejim, (2020), Uzoegwu, and Onwurah (2021), Nwaorgu and Orajika (2016), Okafor and Oko-Ose (2019). These researchers reported higher prevalences in males than females.

From this study, the age range 15-25 had the highest prevalent rate of 17.14% for males and 14.29% for females, followed by 26-35 with prevalent rate of 15.23% for males and 13.33%, while the age range 55-60 had the least prevalent rate of 0.95% for both male and female and this as in conformity with findings of Aribodor *et al.* (2021), Nwaorgu and Orajika (2019) which all

reported decline in prevalence by age.

The mean Glucose values were significantly lower for malaria parasitaemia subjects for both male and female. A significant decline in glucose value with increase in severity was observed for glucose. It was observed from this study that glucose reduction in malaria parasitaemia was not affected by age. Some studies have reported that Plasmodium falciparum parasites fully depend on glucose as an energy source, hence hypoglycemia occurs during the management of patients with malaria (Godyer and Taraschi, 2014), Binh *et al.* (2019) reported hypoglycemia in malaria infected patients, stated host glucose production becomes insufficient for both host and parasite demand as infection progressed. Contrary to this study, Esan (2016) reported that Glucose level in control subjects were observed lower compared to value obtained in pretreatment and post-treatment during malaria parasite infection.

This study showed decreased Total cholesterol and triglyceride levels when compared with control values, though it was not statistically significant, this was in conformity with a systematic review on Serum lipids and lipoproteins in malaria by Benjamin Visser *et al.* (2018). The increased oxidative stress in malaria, which accounts for the degradation of the lipoproteins, may originate from several sources including intracellular of parasitized erythrocytes and extracellular of haemolysed erythrocytes or host immune responses (Memon *et al.*, 2018). These events may be related to the decrease, as significant decrease in total cholesterol levels with increase in severity was observed.

This study established the fact that *P. falciparum* infection has significant effect on the liver, as results of this study showed some significant increases in enzymes activities of aspartate amino transferase (AST), alanine amino transferase (ALT), and alkaline phosphatase (ALP) among patients with *P. falciparum* malaria parasitaemia, which increased with increase in severity of the malaria parasitaemia. The increased serum levels of hepatic enzymes AST and ALT, and ALP are the biomarkers of liver disorders. This results was in conformity with studies by Ignatius *et al.* (2016); Onyesom and Onyemakonor (2017); Mohamed Al-Salahy *et al.* (2016).

The study revealed that total protein, urea, albumin and globulin values of malaria parasitaemia subjects were not statistically significant in comparison with the healthy control, this was at variance with the study done by Melita *et al.* (2021); Prakash *et al.* (2018) who recorded significant levels in malaria infected patients. Significantly higher values were recorded for Creatinine in both males and females, and these values increased significantly with increase in severity. It is well known that creatinine concentration is used for the assessment of renal efficiency, hence when their values are higher than normal, deficiency in renal function is suspected. Elevation of plasma creatinine concentration could be suggestive of ineffective filtering ability of the kidneys which could lead to renal function impairment. This was consistent with the studies by Parkash *et al.* (2018); Ogbadoyi and Tsado (2020).

**Table 5:** Some Biochemical Profile of Plasmodium falciparum Infected inhabitants of Billiri metropolis based on Age.

Parameter	15-25years	26-35 years	36-45 years	46-55 years	55-60 years	f-ratio	p-values
Glucose (mg/dl)	68.12±5.23	66.24±5.66	70.07±6.23	66.43±8.91	67.43±6.87	6.67	1.23 0.68
Total Cholesterol (mg/dl)	146.56±30.12	149.72±24.26	150.23±20.34	147.35±27.18	145.58±23.11	4.93	0.09 0.26
Triglyceride (mg/dl)	80.37±19.24	69.26±12.56	62.34±23.43	78.48±18.34	79.79±13.67	5.07	1.08 0.06
ALT (IU/L)	23.21±2.28	20.14±1.79	21.66±0.79	19.29±2.11	24.16±34	7.12	0.16 0.08
AST (IU/L)	26.34±1.79	25.56±2.52	27.34±3.01	26.67±2.10	22.78±1.65	4.19	0.15 0.06
ALP (mg/dl)	74.17±4.81	69.18±3.78	67.18±8.12	65.84±6.43	70.21±9.21	7.89	1.3
Albumin (g/dl)	3.66±0.51	3.45±0.11	3.54±0.20	3.19±0.18	3.65±0.67	3.02	
Total protein (g/dl)	6.52±0.54	6.25±1.23	6.41±0.97	6.37±0.24	6.74±1.23	8.06	
Globulin (g/dl)	2.86±0.21	2.80±0.43	2.87±0.65	3.18±0.29	3.09±0.48	3.97	
Creatinine (mg/dl)	0.86±0.52	1.03±0.32	1.11±0.37	0.98±0.28	1.02±0.55	6.23	
Urea (mg/dl)	15.08±2.14	18.11±1.45	16.58±2.01	14.98±1.22	16.18±2.31	3.78	

## CONCLUSION

Glucose, Total cholesterol, Liver enzymes, creatinine changes were seen involved in malaria infection caused by *Plasmodium falciparum*. The prevention of malarial infection is still the most valid method of preventing these conditions, early diagnosis and treatment are the measures likely to decrease malarial complications commonly determined by changes in these biochemical parameters in developing countries. Prompt visits to centers equipped to provide necessary anti malarial therapy and support could further reduce mortality and enhance recovery.

## Acknowledgement

The author thanks the Department of Biochemistry Gombe State University and Federal Teaching Hospital Laboratory, for their facility for the research. The Author also appreciates TETFund for sponsorship

## REFERENCES

- Singh, B., Kim Sung and Matusop. A., (2015): A large focus of naturally acquired plasmodium Knowlesi; Infectious in human beings. *Lancet*. 363 (9414): 1017-1024.
- Mueller, I., Zimmerman, P. A., and Reeder, J. C., (2019): Plasmodium malaria and plasmodium ovale in the “bashful” malaria parasitism Trends *Parasitol* 23 (6): 278-283.
- Prothero, R. Mansell E., (2020): Malaria forest and people in southeast Asia-Singapore. *Journal of Tropical Geography*. 20 (1); 76-85.
- Lindemann, M. (2018): Medicine and society in early modern Europe. Cambridge University Press. Pp 62.
- Gratz, N. G., (2017): The vector and rodent borne diseases of Europe and North America. Their distribution and public health burden. Cambridge University Press.
- WHO (2021): Severe plasmodium falciparum malaria; Trans. Royal Soc. Trop. Med. Hug. 95; 51-59.
- Bledsoe, G. H., (2017): Malaria for clinicians in the United State. *South Med. J.* 98 (12); 1197-1204.
- Sturn, A., Amino, R., Van de Sand, C., Regen, T., Retzlaff, S., Rennenberg, A., Krueger A., Pollok, J. M., Menard, R., and Heussler, V. T., (2019): Manipulation of host hepatocytes by the malaria for delivery into liver sinusoids. *Science*. 313 (5791); 1287-1290.
- Cogswell, F. B., (2019): The hypnozoite and relapse in primate malaria. *CliniMicrobio. Rev.* 5 (1); 26-35.
- Angus MGN, Fletcher KA, Maegraith BG. Studies on the lipids of *Plasmodium knowlesi* infected rhesus monkeys (*Macacamulatta*). IV. Changes in erythrocyte lipids. *Ann Trop Med Parasitol*. 1971; 12: 429-439. Advances in Biomedical Sciences 2018; 3(1): 1
- Angus, MGN, Fletcher KA, Maegraith BG. (2015) Studies on the lipids of *Plasmodium knowlesi* infected rhesus monkeys (*Macacamulatta*). II. Changes in serum nonesterified fatty acids. *Ann Trop Med Parasitol*.; 12: 155-167.
- Chagnon A, Guiguen Y, Sutre E. (1985) Hypocholesterolemia in malaria: an aid to diagnosis? *Semaine des Hopitaux*. 1985; 12: 2075-2076.
- National Population Commission (2006): Housing and population census result: Gombe State National population Office, Benin City.
- NIPOST (2009): Post Offices-with map of LGA. Archived from the

- original on 2009-10-07. Retrieved 2009-10-20.
- Dayachi, F., Kabongo, L. and Ngoie, K. (2018): Decreased mortality from Malaria in children with symptomatic HIV infection. *Int. Cont. AIDS*. 2: 164.
- Monica Cheesbrough. (2005). Discrete Laboratory Practice in Tropical Countries Part 1, Cambridge Second Editions. Published by Press Syndicate of the University of Cambridge, chp. 5, page 247-258.
- Warhurst, D. C. and Williams, J. E. (1996): Acp Broadsheet no 148. July 1996. Laboratory diagnosis of malaria. *J. Clin. Pathol.* 49: 533-538.
- Reitman, S. and Frankel, S. (1957): A colorimetric method for the determination of Aspartate amino transferase and Alanine amino transferase activities in serum. *Amer. J Clin Path*; 28: 56.
- Rec. Gscc (DGKC) (1972). Optimised Standard Colorimetric Methods. *J. Clin. Chem. Clin. Biochem.*, 10: 182.
- Spencer, K. and Price, C. P. (1977). Simple screening tests for serum albumin levels. *Annals of Clinical Biochemistry*, 14: 105-115.
- Slot, C. (1965). Blood creatinine level an renal function. *Journal of Clinical Laboratory Investigation*. 17: 891-895.
- Kaplan, A. (1965). Standard Methods of Clinical Chemistry. New York, Academic Press pp. 256.
- Rey and Wielinger (1970). Glucose oxidase method. *Z. analyt. chem.* 252: 224.
- Allain, C. C., Poon, L. S., Chan, C. S. G. (1974): Enzymatic determination of total serum cholesterol. *Clinical Chemistry*. 20, 470.
- Mbanugo, J. I. and Ejims, D. O. (2020). Plasmodium Infections in Children Aged 0-5 yrs in Awka Metropolis, Anambra State, Nigeria. *Nigeria Journal of Parasitology*. 2: 55-59.
- Uzoegwu, P. N. and Onwurah, A. E. (2021). correlation of lipid Peroxide in malariapositive and negative status of AA, AS and SS individuals from the University of Nigeria Nsukka community. *Journal of Bio Research and Biotechnology*, 1 (1): 97-114.
- Nwaorgu, O. C. and Orajaka, B. N. (2016). Prevalence of malaria among Children 1-10 years old in Communities in Awka North Local Government Area, Anambra State South East Nigeria. *International Multidisciplinary Journal Ethiopia*, 5 (5): 264-281.
- Okafor, F. U. and Oko-Ose, J. N. (2019). Prevalence of malaria infections among children aged six months to eleven years (6 months-11 years) in a tertiary institution in Benin City, Nigeria. *Global Advanced Journal and Medical Sciences*, 1 (10): 273-279.
- Aribodor, D. N., Njoku, O. O., Eneanya, C. I. and Onyali, I. O. (2020). Studies on Prevalence of Malaria and management practices of the Azia Community in Ihiala L. G. A., Anambra State, South-East Nigeria. *Nigeria Journal of Parasitology*, 22: 42-48.
- Goodyer, I. D., Taraschi, T. F. (2014). Plasmodium falciparum: a simple, rapid method for detecting parasite clones in microtiter plates. *Exp. Parasitol.*, 86: 158-160.
- Davis, T. M., Binh, T. Q., Thu le, T. A., Long, T. T., Johnston, W., Robertson, K. and Barrett, P. H. (2015). Glucose and lactate turnover in adults with falciparum malaria: effect of complications and antimalarial therapy. *Trans. R.*



- Soc. Trop. Med. Hyg.* 96: 411-417.
- Binh, T. Q., Davis, T. M., Johnston, W., Thu, L. T., Boston, R., Danh, P. T. and Anh, T. K. (2019). Glucose metabolism in severe malaria: minimal model analysis of the intravenous glucose tolerance test incorporating a stable glucose label. *Metabolism*. 46 (12): 1435-1440.
- Esan, A. J. (2016): Incidence and Evaluation of Stress Induced In Plasmodium Falciparum Malaria Infected Individuals Using Cortisol, Malondialdehyde, Blood Glucose and Lipid Profile Level. *Journal of Medical and Biological Science Research* Vol. 2 (10), pp. 158-162, December, 2016 ISSN: 2449-1810.
- Benjamin J. Visser, Rosanne W. Wieten, Ingeborg M. Nagel, and Martin P. Grobusc (2018): Serum lipids and lipoproteins in malaria - a systematic review and meta-analysis. *Malar J.* 12: 442.
- Memon, R. A., Staprans, I., Noor, M., Holleran, W. M., Uchida, Y., Moser, A. H., Feingold, K. R., Grunfeld, C. (2018). Infection and inflammation induce LDL oxidation in vivo. *Arterioscler. Thromb. Vasc. Biol.* 20: 1536-1542.
- Ignatius, C. M., Emeka, E. N. and Blessing, N. E. (2016): "Effect of malaria parasitaemia on liver enzyme tests," *International Journal of Tropical Medicine*, vol. 3, no. 3, pp. 49-52.
- Onyesom, I. and Onyemakonor, N. (2017): "Levels of parasitaemia and changes in some liver enzymes among malarial infected patients in Edo-Delta Region of Nigeria," *Current Research Journal of Biological Sciences*, vol. 3, no. 2, pp. 78-81.
- Mohamed Al-Salahy, Bushra Shnawa, GamalAbed, Ahmed Mandour, and Ali Al-Ezzi. (2016): Parasitaemia and Its Relation to Hematological Parameters and Liver Function among Patients Malaria in Abs, Hajjah, Northwest Yemen. *Interdisciplinary Perspectives on Infectious Diseases*; Volume 2016, Article ID 5954394, 5 pages.
- Melita, K. S., Halankar, A. R., Makwana, P. D., Torane, P. P., Satija, P. S. and Shah, V. B. (2021). Severe acute renal failure in Malaria. *Journal of Postgraduate Medicine*, 47 (1): 24-26.
- Parkash, J., Singh, A. K., Kumar, N. S. and Saxena, R. K. (2018). Acute renal failure in Plasmodium vivax malaria. *Journal of Association of Physicians, India*. 51: 265-267.
- Ogbadoyi, E. O and Tsado, R. D. (2020). Renal and Hepatic Dysfunction in Malaria patients in Minna, North Central Nigeria. *Online Journal of Health and Allied Sciences*, 8 (3): 3-8.